

RESEARCH NOTE

The link between mother and adolescent substance use: inter-generational findings from the British Cohort Study

Megan E. Patrick, University of Michigan, USA

meganpat@umich.edu

Jennifer L. Maggs, Pennsylvania State University, USA

Kaylin M. Greene, Montana State University, USA

Nicole R. Morgan, Pennsylvania State University, USA

John E. Schulenberg, University of Michigan, USA

(Received January 2013 Revised August 2013)

Abstract

The objective of this study was to identify mother, family, and individual factors associated with adolescent alcohol, tobacco, and marijuana use, using mother and child self-reports. Adolescents aged 12-15 (N=276) and their mothers, who were participants in the British Cohort Study (BCS; born 1970), were both surveyed when mothers were 34 years old. Predictors included mothers' substance use as well as characteristics of the child (gender, age, conduct problems) and family (social class, two-parent family, parent-adolescent conflict). Outcome variables were adolescent alcohol, cigarette, and marijuana use. Child characteristics were predictive, with older children more likely to engage in all behaviors. After controlling for other predictors, mothers' current drinking frequency and problems predicted adolescent reports of ever drinking and of drinking sometimes/regularly; mothers' marijuana use was a marginally significant predictor of adolescent marijuana use. Results suggest that mothers' substance use is an important component of adolescent use, even after accounting for characteristics of the child and the intergenerational family context.

Key words: parental substance use; mothers; adolescents; alcohol; drinking; marijuana/cannabis; tobacco

Introduction

A large body of literature has explored family factors that may contribute to the inter-generational transmission of substance use. This research suggests that families can shape adolescent alcohol and drug use through a number of pathways including parental human capital resources, parent-child interactions and relationships, and behavior modeling (Vakalahi, 2001). Positive family relationships, such as parent-child closeness, are linked to less substance use whereas

parent-child conflict is associated with greater substance use (Hawkins, Catalano, & Miller, 1992; Kuntsche & Silbereisen, 2004). In addition, parental educational and other resources have been linked to youth substance use (e.g., Koning, Engels, & Verdurmen, 2010; Maggs, Patrick, & Feinstein, 2008; Melotti et al., 2011). Parental modeling and other inter-generational transmission mechanisms are also important, given the large body of research suggesting that parents who engage in heavy drinking and use drugs are more likely to have children who drink alcohol and use drugs

themselves (Donovan et al., 2004; Dooley & Prause, 2007; Fawzy, 1983; Osborne & Berger, 2009; Zucker, Jester, Fitzgerald, Puttler, & Wong, 2003).

Although associations between parent and child substance use have been well-documented, extant research is unclear about whether this association remains after accounting for confounding factors that may be correlated with both parental and child substance use. In other words, it is uncertain whether parental substance use itself matters (e.g., having available substances or modeling alcohol and drug use) or whether other factors—such as socio-economic status or family relations—account for this association. Whereas many studies continue to find a robust association after controlling for a host of family and individual characteristics (e.g., Hemphill et al., 2011), others find that accounting for these factors reduces the association between parent and child substance use to non-significance (e.g., Koning et al., 2010). In addition, a link between parent and youth substance use has been found for some substances, but not others, once control variables are included in the models (Macleod et al., 2008; Schinke, Fang, & Cole, 2008).

The present study builds on and advances this literature by testing the extent to which mothers' substance use is associated with the substance use of their adolescent children, using self-reports from both generations. We add to the literature in three primary ways. First, we control for a number of family factors, to better understand whether the link between parental and youth substance use remains after accounting for factors predictive of substance use for both generations. Second, we test the inter-generational linkages across the three most commonly-used substances (i.e., alcohol, cigarettes, and marijuana) rather than focusing on only one type of substance use. Third, we utilize self-report measures of substance use from both mothers and adolescents, which enables us to get more accurate information from each generation and reduces single-reporter bias. Understanding predictors of early substance use is critical in the light of research demonstrating that early onset of alcohol and drug use predicts substance use and disorders in late adolescence and young adulthood (Agrawal et al., 2006; Clark, Cornelius, Kirisci, & Tarter, 2005; Dooley, Prause, Ham-Rowbottom, & Emptage, 2005; Grant and Dawson, 1997).

Method

Data are from the British Cohort Study (BCS), an ongoing birth cohort study that sampled all individuals in Britain who were born in one week in 1970 (Butler, Golding, & Howlett, 1985; Bynner, Butler, Ferri, Shepherd, & Smith, 2000; Schoon, 2006). After an initial assessment of 16,571 infants (96% of live births), follow-ups were conducted at ages 5, 10, 16, 26, 30, 34, 38, and 42. Multiple sources of data were collected. Parent data in the present analyses were obtained from the primary BCS cohort members in 2004, when they were age 34 ($N=9316$; $n=5039$ women; 70% retention excluding emigrants and deceased). At age 34, 15% ($N=772$) of BCS women had at least one biological or adopted child aged 12 to 15 years old (born 1989-1992) who was living with them¹, 49% of whom ($N=375$) were selected for a sub-study about parents and children. Of those selected, 36 were missing all parent or child data, 60 were missing data on covariates of interest, and 3 lacked a parent report and child self-report for the same child, yielding a final analytic sample size of 276 (74% of those eligible). Data comprised mother reports (interview and self-completion) of demographics, parent-child conflict, and child temperament; and adolescent reports (self-completion) of their own attitudes and behaviors (see also Simmonds, Fuller, Lessof, & Foudouli, 2007). Only the eldest child of each mother was included.

Measures

Adolescent substance use outcomes

Adolescents were asked how often they used alcohol, cigarettes, and marijuana, with response options of (a) I have never drunk alcohol; (b) I have drunk alcohol once or twice only; (c) I used to drink alcohol but I don't now; (d) I sometimes drink alcohol, but I don't drink alcohol every week; and (e) I drink alcohol regularly, once a week or more. This measure of drinking frequency was used to derive primary dichotomous dependent variables in which youth who self-reported that they *ever drank alcohol* (i.e., responded b-e, coded 1) were compared those who never drank alcohol (i.e., responded a, coded 0). Those who *drank alcohol sometimes or regularly* (i.e., d-e, coded 1) were compared to those who drank less often (i.e., a-c, coded 0). Other substances were coded using the same response scale as alcohol. Adolescents who

reported *ever using cigarettes* or *ever using marijuana* (coded 1) were compared to those who had never used the substance (coded 0).

Mothers' substance use

Mothers' current drinking frequency (i.e., how often she has an alcoholic drink of any kind) was reported on a scale of 0=never had an alcoholic drink, 1=never nowadays, 2=less often or only on special occasions, 3=two to three times a month, 4=once a week, 5=two to three days a week, 6=on most days. *Mothers' problem drinking* was measured with the CAGE (Bradley, Boyd-Wickizer, Powell, & Burman, 1998; Ewing, 1984; Liskow, Campbell, Nickel, & Powell, 1995; Mayfield, McLeod, & Hall, 1974), which asked mothers whether they had ever experienced 4 alcohol-related symptoms in their lifetime and, if so, whether the symptoms had occurred in the past year. Responses were coded to contrast those who reported one or more alcohol problems in the prior year (coded 1) with those reporting no problems in the prior year (coded 0). *Mothers' frequency of cigarette use* was coded as 0=has never smoked cigarettes, 1=used to smoke cigarettes but does not now, 2=smokes cigarettes occasionally but not every day, 3=smokes cigarettes every day. *Mothers' frequency of marijuana use* was coded as 0=has never tried cannabis, 1=never nowadays, 2=less often or only on special occasions, 3=once a month, 4=two or three times a month, 6=two or three days a week, 7=most days.

Adolescent characteristics

Adolescent controls were *gender* (male [1] vs. female [0]), *age* (14-15 years [1] vs. 12-13 years [0]), and *conduct problems*, measured with the conduct problems subscale of the Strengths and Difficulties questionnaire ($\alpha=.75$; Goodman, 1997) as reported by the mothers. The scale is the sum of five items with response options ranging from 0=not true to 2=certainly true.

Family characteristics

Family social class/ employment was coded based on two measures. First, the Registrar General's social class measure, a categorization of six social groups based on current occupational attainment, was used to classify mothers and their partners into skilled versus unskilled/partly skilled occupations (Rose & Pevalin,

2001). The parent/guardian with the highest occupational status was used to determine the social class of the family, as non-manual (including professional/managerial) status or manual status. In addition, a dichotomous variable captured if the mother was employed or not employed (i.e., unemployed, disabled, or looking after the family). The two variables (social class and mother employment) were coded together into three groups: non-manual social class with mother employed [reference group], manual social class, and mother not employed. Living situation was included with adolescents living with *two parents* coded as 1 (all else=0). Last, *parent-adolescent conflict* was assessed with the conflict scale of the Child-Parent Relationship Scale-Short Form (Pianta, 1992). The scale was created by summing the 7 items ($\alpha=.86$; e.g., "My child easily becomes angry at me"), with response options for each item ranging from 0=definitely does not apply to 4=definitely applies to me.

Results

Descriptive statistics regarding predictor and outcome variables are shown in Table 1. Although the current sample of adolescents is not nationally representative, the substance use patterns are generally consistent with those of the UK adolescent population as a whole.ⁱⁱ Mothers of young adolescents included in the analyses presented here were similar to other women in the BCS 1970 cohort: there were no significant differences for marijuana use or CAGE scores. However, mothers of young adolescents were more likely to be daily cigarette smokers than other women. Logistic regression analyses were used to predict adolescent substance use as a function of adolescent gender, age, conduct problems, family social class, mothers' employment, two-parent family status, parent-adolescent conflict, and mothers' substance use. Indicators of mothers' substance use were tested in separate models, due to collinearity between the two indicators of alcohol use and problems, and our interest in testing domain-specific transmission of substance use. Results shown are multi-variate, due to our primary interest in whether the link between maternal and youth substance use remained after accounting for other individual and family factors.

Table 1. Descriptive statistics for British Cohort Study mothers (age 34) and their adolescent children (age 12-15) (N=276)

	% or M (SD)
Adolescent characteristics	
Male gender	52.2%
Older age (14-15 years)	44.6%
Conduct problems	2.00 (2.05)
Ever drank alcohol	76.4%
Sometimes drank alcohol	30.8%
Ever smoked cigarettes	34.4%
Ever used marijuana	10.2%
Family characteristics	
Social class - non-manual	49.3%
Social class - manual	33.7%
Mother not employed	29.7%
Two-parent family	72.5%
Parent-adolescent conflict	9.21 (6.88)
Mother characteristics	
Mothers' current drinking frequency	4.3% Never 7.2% Never nowadays 23.2% Less often or only on special occasions 15.9% 2-3 times/month 19.2% Once a week 21.4% 2-3 days/week 8.7% Most days
If drinks, units of alcohol in prior week	6.31 (9.12)
Mothers' drinking problems	20.6% ^a
Mothers' smoking frequency	29.0% Never 21.4% Used to smoke, but not at all now 7.2% Occasionally, but not every day 42.4% Every day
If smokes, number of cigarettes per day	15.75 (7.30)
Mother current marijuana use frequency	82.5% Never tried 5.5% Never nowadays 7.3% Less often or only on special occasions 0.7% Once a month 0.7% Two to three times a month 0.7% Two to three days a week 2.6% On most days

Note: ^aThere were 14 missing cases for the mothers' CAGE score.

Table 2. Logistic regressions predicting adolescent substance use by adolescent, family, and mother characteristics

	Ever drank alcohol N=262 OR [CI]	Sometimes drank alcohol N=262 OR [CI]	Ever smoked cigarettes N=276 OR [CI]	Ever used marijuana N=273 OR [CI]
Intercept	1.05	0.11 ^{***}	0.13 ^{***}	0.00 ^{***}
Mother characteristics				
Mothers' drinking problems	2.50 [1.01 6.16] [*]	2.11 [1.05, 4.26] [*]	---	---
Mothers' smoking frequency	---	---	1.10 [0.87, 1.39]	---
Mothers' frequency of marijuana use	---	---	---	1.32 [0.99, 1.76] ⁺
Adolescent characteristics				
Male gender	0.73 [0.38, 1.41]	1.01 [0.56, 1.84]	0.43 [0.24, 0.78] ^{**}	1.92 [0.75, 4.93]
Older age (14-15 years)	5.34 [2.51, 11.38] ^{***}	6.38 [3.49, 11.66] ^{***}	5.52 [3.08, 9.87] ^{***}	43.72 [5.61, 340.48] ^{***}
Conduct problems	1.20 [0.92, 1.55] ⁺	1.05 [0.86, 1.29]	1.21 [0.99, 1.47] ⁺	1.20 [0.93, 1.54] ⁺
Family characteristics				
Social class ^a				
Manual social class (unskilled/partly skilled)	0.73 [0.38, 1.41]	0.76 [0.40, 1.43]	0.84 [0.45, 1.57]	1.25 [0.49, 3.17]
Mother not employed	0.97 [0.49, 1.94]	1.05 [0.55, 2.00]	1.08 [0.57, 2.02]	1.04 [0.39, 2.80]
Two-parent family	1.83 [0.91, 3.68] ⁺	0.96 [0.50, 1.84]	1.15 [0.61, 2.20]	1.01 [0.37, 2.75]
Parent-adolescent conflict	1.01 [0.94, 1.08]	1.03 [0.97, 1.09]	1.04 [0.98, 1.10]	1.04 [0.95, 1.13]

^aReference group: one or both parents reported professional/managerial/technical/skilled non-manual social class, and the mother was employed.

⁺ $p < .10$ (one-tailed test, $p < .05$), ^{*} $p < .05$, ^{**} $p < .01$, ^{***} $p < .001$. OR = odds ratios. CI = 95% confidence intervals.

With regards to predicting adolescent drinking (see Table 2), while controlling for the adolescent and family characteristics, adolescents whose mothers reported at least one alcohol problem in the prior year, as indexed by the CAGE, had greater odds of ever and of sometimes drinking. In these multivariate models, none of the other adolescent or family predictors was significant, with the exception of age, with 14-15 year old adolescents showing a much greater likelihood of both ever drinking and of sometimes drinking than 12-13 year old adolescents. Adolescents with more conduct problems had marginally significant greater odds of ever drinking ($p < .10$) and adolescents in two-parent families had marginally significant greater odds of ever drinking ($p < .10$). In additional models (not tabled), adolescents whose mothers drank more frequently, also evidenced greater odds of ever drinking (OR=1.44, CI=[1.18, 1.76], $p < .001$) and of sometimes drinking (OR=1.39, CI=[1.15, 1.69], $p < .001$).

In terms of predicting adolescents' likelihood of ever smoking cigarettes, while controlling for adolescent and family characteristics, mothers' smoking did not predict the odds of adolescent smoking. In these multivariate models, none of the family predictors was significant, but the adolescent predictors were: boys were less likely, and 14-15 year olds were more likely, to have smoked. Conduct problems approached significance ($p < .10$) as a positive predictor of ever using cigarettes.

Finally, in reference to predicting the likelihood of adolescents ever having used marijuana, while controlling for adolescent and family characteristics, mothers' marijuana use was a marginally significant predictor of a greater likelihood of adolescent marijuana use ($p < .10$). In a separate model (not tabled), the frequency of mothers' current marijuana use was a marginally significant predictor of adolescent marijuana use (OR=1.32, CI=[0.99, 1.76], $p < .10$). None of the family predictors was significant. The relatively older adolescents were more likely to have used marijuana. Conduct problems were marginally significant as a positive predictor of ever using marijuana ($p < .10$).

Discussion

There is little doubt that as a psychosocial system the family contributes extensively to adolescent substance use (Hawkins et al., 1992; Kuntsche & Silbereisen, 2004; Vakalahi, 2001). However, adequately specifying the inter-generational links between substance use and abuse by mothers and children remains difficult (Hemphill et al., 2011; Koning et al., 2010). This study addresses some key gaps in the literature by including several possible family factors, multiple forms of adolescent substance use, and both mothers' and children's reports. Our key findings are that, after controlling for other individual and family factors, mothers' current drinking problems predicted adolescent drinking. In addition, mothers' current marijuana use approached significance predicting adolescent marijuana use. These findings are in line with other research highlighting linkages between maternal and child substance use (e.g. Dooley & Prause, 2007; Macleod et al., 2008).

It is notable that maternal substance use remained among the strongest predictors of adolescent substance use even after controlling for other family and child effects. Prior theory and research suggest that mothers may shape adolescent substance use in various ways. Mothers may model substance use for their children and they may shape youths' access to substances (e.g., by having their preferred substances accessible at home). Mothers may also influence social norms and perceptions of the risks and benefits of early substance use. Furthermore, problematic maternal substance use may increase adolescent substance use by interfering with successful parenting strategies such as adolescent monitoring. It is also possible that the associations documented in the current study result not from environmental factors but from genetic similarity between mother and child (Rowe, 1994). The current study does not test these mediating pathways, and thus cannot identify which may be responsible for the inter-generational transmission of substance use. However, regardless of the underlying pathways, the results of the current study suggest that policies and interventions should aim to identify youth whose mothers are problematic substance users. These youth are particularly at risk for substance use during early adolescence and thus

early intervention among this population may be especially important.

This national sample of mothers born in 1970 brings important strengths to the understanding of family and child influences on adolescent substance use. Large national data sets offer confidence in terms of population coverage and generalizability. However, the sampling frame included the mothers when they themselves were children, so the sample is potentially biased due to any selective attrition. The relatively small sample size also leads to a lack of power to detect statistical significance for some findings. In addition, the sample is restricted to mother-adolescent dyads in which the children were born when mothers were aged 19 to 22 (in 1989 to

1992); therefore, it is unclear whether the results would generalize to mothers who were older when their children were born. Of course, this age limitation also is advantageous because it reduces the age heterogeneity that typically arises for the second generation, when the first generation is the sampling target. Strengths of the study include that having both maternal and child reports helps overcome limitations of sole-source data. Future research, building on our findings and using a larger sample, should test mediational models specifying family mechanisms that connect maternal and adolescent substance use and abuse, as well as the family moderators of this link.

Acknowledgements

Funding for data analyses was provided by the National Institute on Alcohol Abuse and Alcoholism (R01 AA019606 to J. Maggs). We are grateful to The Centre for Longitudinal Studies, Institute of Education for the use of these data and to the UK Data Archive and Economic and Social Data Service for making them available. However, they bear no responsibility for the analysis or interpretation of these data.

References

- Agrawal, A., Grant, J. D., Waldron, M., Duncan, A. E., Scherrer, J. F., Lynskey, M. T., Madden, P. A., Bucholz, K. K., & Heath, A. C. (2006). Risk for initiation of substance use as a function of age of onset of cigarette, alcohol and cannabis use: Findings in a Midwestern female twin cohort. *Prevention Medicine*, 43, 125-128. <http://dx.doi.org/10.1016/j.ypmed.2006.03.022>
- Bradley, K. A., Boyd-Wickizer, J., Powell, S. H., & Burman, M. L. (1998). Alcohol screening questionnaires in women: A critical review. *The Journal of the American Medical Association*, 280, 166-171. <http://dx.doi.org/10.1001/jama.280.2.166>
- Butler, N. R., Golding, J., & Howlett, B. C. (1985). *From birth to five: A study of the health and behaviour of a national cohort*. Oxford: Pergamon.
- Bynner, J., Butler, N., Ferri, E., Shepherd, P., & Smith, K. (2000). The design and conduct of the 1999-2000 surveys of the National Child Development Study and the 1970 British Cohort Study (CLS Cohort Studies Working Paper 1). London: University of London, Institute of Education, Centre for Longitudinal Studies.
- Clark, D. B., Cornelius, J. R., Kirisci, L., & Tarter, R. E. (2005). Childhood risk categories for adolescent substance involvement: A general liability typology. *Drug and Alcohol Dependence*, 77, 13-21. <http://dx.doi.org/10.1016/j.drugalcdep.2004.06.008>
- Donovan, J. E., Leech, S. L., Zucker, R. A., Loveland-Cherry, C. J., Jester, J. M., Fitzgerald, H. E., Putter, L. E., Wong, M. M., & Looman, W. S. (2004). Really underage drinkers: Alcohol use among elementary students. *Alcohol: Clinical & Experimental Research*, 28, 341-349. <http://dx.doi.org/10.1097/01.ALC.0000113922.77569.4E>
- Dooley, D., & Prause, J. (2007). Predictors of early alcohol drinking onset. *Journal of Child & Adolescent Substance Abuse*, 16, 1-29. http://dx.doi.org/10.1300/J029v16n02_01
- Dooley, D., Prause, J., Ham-Rowbottom, K. A., & Emptage, N. (2005). Age of alcohol drinking onset: Precursors and the mediation of alcohol disorder. *Journal of Child & Adolescent Substance Abuse*, 15, 19-37. http://dx.doi.org/10.1300/J029v15n02_02
- Ewing, J. A. (1984). Detecting alcoholism: The CAGE questionnaire. *The Journal of the American Medical Association*, 252, 1905-1907. <http://dx.doi.org/10.1001/jama.1984.03350140051025>
- Fawzy, F. I. (1983). Generational continuity in the use of substances: The impact of parental substance use on adolescent substance use. *Addictive Behaviors*, 8, 109-114. [http://dx.doi.org/10.1016/0306-4603\(83\)90003-5](http://dx.doi.org/10.1016/0306-4603(83)90003-5)
- Goodman, R. (1997). Strengths and difficulties questionnaire: A research note. *The Journal of Child Psychology and Psychiatry*, 38, 581-586. <http://dx.doi.org/10.1111/j.1469-7610.1997.tb01545.x>
- Grant, B. F., & Dawson, D. A. (1997). Age at onset of alcohol use and its association with DSM-IV alcohol abuse and dependence: Results from the National Longitudinal Alcohol Epidemiologic Survey. *Journal of Substance Abuse*, 9, 103-110. [http://dx.doi.org/10.1016/S0899-3289\(97\)90009-2](http://dx.doi.org/10.1016/S0899-3289(97)90009-2)

- Hawkins, J. D., Catalano, R., & Miller, J. (1992). Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: Implications for substance abuse prevention. *Psychology Bulletin*, 112, 64-105.
<http://dx.doi.org/10.1037/0033-2909.112.1.64>
- Hemphill, S. A., Heerde, J. A., Herrenkohl, T. I., Patton, G. C., Toumbourou, J. W., & Catalano, R. F. (2011). Risk and protective factors for adolescent substance use in Washington state, the United States and Victoria, Australia: A longitudinal study. *Journal of Adolescent Health*, 49, 312-320. <http://dx.doi.org/10.1016/j.jadohealth.2010.12.017>
- Hibell, B., Andersson, B., Bjarnason, T., Ahlström, S., Balakireva, O., & Kokkevi, A. (2004). The ESPAD report 2003. Alcohol and other drug use among students in 35 European countries. Stockholm, Sweden: The Swedish Council for Information on Alcohol and Other Drugs-The Pompidou Group at the Council of Europe. Downloaded on July 26, 2013 from <http://www.espad.org/en/Reports--Documents/ESPAD-Reports/>
- Koning, I.M., Engels, R. C. M. E., & Verdurmen, E. E. (2010). Alcohol-specific socialization practices and alcohol use in Dutch early adolescents. *Journal of Adolescence*, 33, 93-100. <http://dx.doi.org/10.1016/j.adolescence.2009.05.003>
- Kuntsche, E. N., & Silbereisen, R. K. (2004). Parental closeness and adolescent substance use in single and two-parent families in Switzerland. *Swiss Journal of Psychology*, 63, 85-92. <http://dx.doi.org/10.1024/1421-0185.63.2.85>
- Liskow, B., Campbell, J., Nickel, E., & Powell, B. J. (1995). Validity of the CAGE questionnaire in screening for alcohol dependence in a walk-in (triage) clinic. *Journal of Studies on Alcohol and Drugs*, 56, 277-281.
- Macleod, J., Hickman, M., Bowen, E., Alati, R., Tilling, K., & Smith, G. D. (2008). Parental drug use, early adversities, later childhood problems and children's use of tobacco and alcohol at age 10: Birth cohort study. *Addiction*, 103, 1731-1743.
<http://dx.doi.org/10.1111/j.1360-0443.2008.02301.x>
- Maggs, J. L., Patrick, M. E., & Feinstein, L. (2008). Childhood and adolescent predictors of alcohol use and problems in the National Child Development Study. *Addiction*, 103 (Suppl. 1), 7-22. <http://dx.doi.org/10.1111/j.1360-0443.2008.02173.x>
- Mayfield, D., McLeod, G., & Hall, P. (1974). The CAGE questionnaire: Validation of a new alcoholism screening instrument. *The American Journal of Psychiatry*, 131, 1121-1123.
- Melotti, R., Heron, J., Hickman, M., Macleod, J., Araya, R., & Lewis, G. (2011). Adolescent alcohol and tobacco use and early socioeconomic position: The ALSPAC birth cohort. *Journal of Pediatrics*, 127, e948-955.
<http://dx.doi.org/10.1542/peds.2009-3450>
- Osborne, C., & Berger, L. (2009). Parental substance abuse and child well-being: A consideration of parents' gender and coresidence. *Journal of Family Issues*, 30, 341-370. <http://dx.doi.org/10.1177/0192513X08326225>
- Pianta, R. C. (1992). Child-parent relationship scale. Unpublished measure. Charlottesville, VA: University of Virginia.
- Rose, D., & Pevalin, D. (2001). The national statistics socio-economic classification: Unifying official and sociological approaches to the conceptualisation and measurement of social class. (ISER Working Paper 2001-4). Colchester, England: University of Essex.
- Rowe, D. C. (1994). Genetic and cultural explanations of adolescent risk taking and problem behavior. In R. D. Ketterlinus & M. E. Lamb (Eds.), *Adolescent problem behaviors: Issues and research* (109-126). Hillsdale, NJ: Lawrence Erlbaum.
- Schinke, S. P., Fang, L., & Cole, K. C. A. (2008). Substance use among early adolescent girls: Risk and protective factors. *Journal of Adolescent Health*, 43, 191-194. <http://dx.doi.org/10.1016/j.jadohealth.2007.12.014>
- Schoon, I. (2006). *Risk and resilience: Adaptations in changing times*. Cambridge, England: Cambridge University Press.
<http://dx.doi.org/10.1017/CBO9780511490132>
- Simmonds, N., Fuller, E., Lessof, C., & Foudouli, V. (2007). 1970 British Cohort Study: 2004-2005 Survey Technical Report. Prepared for Centre for Longitudinal Studies, Institute of Education. Accessed online on July 15, 2013 at: <http://www.cls.ioe.ac.uk/page.aspx?&sitesectionid=827&sitesectiontitle=Technical+report>
- Vakalahi, H. F. (2001). Adolescent substance use and family-based risk and protective factors: A literature review. *Journal of Drug Education*, 31, 29-46. <http://dx.doi.org/10.2190/QP75-P9AR-NUVJ-FJCB>
- Zucker, R. A., Jester, J. M., Fitzgerald, H. E., Puttler, L. I., & Wong, M.M. (2003). Predicting onset of drinking from behavior at three years of age: Influence of early child expectancies and parental alcohol involvement upon early first use. *Alcoholism: Clinical & Experimental Research*, 27(Suppl. 5), 174A.

Endnotes

ⁱ Small number of A resident fathers ($n=81$) had children aged 12 to 15. Analyses focused only on mothers due to the smaller sample of fathers. Mother and father data were not combined due to potential differences in the impact of mother and father substance use.

ⁱⁱ National estimates from 2003 documented that 94% of UK youth aged 15-16 had consumed alcohol, 58% had smoked cigarettes, and 38% had used marijuana or hashish. Substance use initiation often occurred early in life: At age 13 (or younger) 61% of UK youth had already consumed at least one glass of beer, 65% had consumed at least one glass of wine, 41% had used cigarettes at least once, and 13% had used marijuana or hashish (Hibell et al., 2004).